



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

OCTOBER 31.

The President, Dr. RUSCHENBERGER, in the chair.

Thirty-five members present.

A paper entitled "Descriptions of Vertebrate Remains chiefly from the Ashley Phosphate Beds of North Carolina," by Jos. Leidy, M.D., was presented for publication.

Self-fertilization in Mentzelia ornata.—Mr. THOMAS MEEHAN referred to an objection made during his remarks on this plant some weeks ago, that a flower which had produced a perfect capsule under a gauze bag to exclude insects, might yet not produce perfect seeds. The capsule was now ripe, and the seed perfect.

Direct Growth Force in Roots.—Mr. MEEHAN spoke of the direct growth-force in roots, as illustrated by some specimens of the White Hickory (*Carya tomentosa*) exhibited. The Hickories during the first few years of their growth developed far more beneath than above the surface. He had seen Pecan Nuts (*Carya olivæformis*), with weak stems not two feet high, have tap roots six feet long. In the one year hickory now exhibited, the tap root was three times the length of the stem. In one plant, however, the young radicle, instead of pushing through the cleft made by the separated shell, had been directed into the shell, and in its fruitless effort to penetrate the wall, had lingered so long, that the upper portion had grown so large as to prevent egress. The root, therefore, instead of making a slender growth of eighteen inches long, had simply made a bulb of about three quarters of an inch in diameter with the shell of the nut attached to it.

Interpretation of varying Forms.—Mr. THOS. MEEHAN said that William Bartram, in the last century, had found forms of *Liriodendron tulipifera* on the Schuylkill River, as he had been informed by his son-in-law, with entire leaves, but only this year had he succeeded in discovering them. Some of these leaves he exhibited. He observed that years ago, such discoveries had an interest in themselves. Now the botanist expected to find entire leaved forms among kinds usually lobed, or lobed ones among the entire leaved class; the only value now in these discoveries is in any lesson they might teach. As a rule, he hesitated to refer to the unpublished observations of others, preferring that the discoverers should in their own good time and way, report what they had found; but hoped to be pardoned on this occasion, for saying that on a recent visit to the Academy, the distinguished botanist Dr. Engel-

leaves, or at least more or less so in comparison with those which they afterwards assume. These large trees with entire leaves had merely retained their juvenescent form. The other attendant characters of juvenescence were also present. The tree from which the large entire leaf exhibited was taken had no signs of ever having borne seeds. In one place he found two trees which, from surrounding circumstances, he should judge were probably about the same age, and in every circumstance relating to nutrition equally favored, one with very deeply cut leaves even to the most feeble branch was covered with seed cones, and was thirteen feet in circumference. The other had leaves almost entire, with but few fruit, and a trunk of only eight feet round.

The danger was that in discussing laws of variation in connection with the origin of species we may overlook these sexual and physiological changes. If one never having seen a Baltimore oriole should notice particularly the brilliant plumage of the male bird, and, without noticing the sex, compare it with the very different looking female bird, he would be very apt to think he had found a "missing link" in a grand evolutionary chain. There were many differences in animals which were recognized as having their origin in obscure sexual laws, as well as many more unrecognized but probable; and he believed these cases were far more numerous in vegetation, and they would have to be carefully eliminated from consideration in any study on the origin of species or the evolution of form in relation thereto.

Edwin A. Barber, H. F. Whitman, and Dr. W. Forwood, U. S. A., were elected members.

Col. W. L. Ludlow was elected a correspondent.

The following papers were ordered to be printed:—